

Precocious Initiation into Smoking, Alcohol Use, and Gambling among Children with Conduct Problems

The Canadian Journal of Psychiatry /
La Revue Canadienne de Psychiatrie
2016, Vol. 61(1) 50-58
© The Author(s) 2016
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0706743715620402
TheCJP.ca | LaRCP.ca



Début précoce du tabagisme, de la consommation d'alcool et du jeu chez les enfants ayant des problèmes de conduite

Caroline E. Temcheff, PhD¹, Michèle Déry, PhD¹,
Renée A. St-Pierre, PhD¹, Myriam Laventure, PhD¹,
and Jean-Pascal Lemelin, PhD¹

Abstract

Objective: Adolescent participation in risky and addictive behaviours, such as smoking, substance use, and gambling has the potential to lead to many serious problems. The presence of conduct problems (CPs) and early initiation into risky and addictive behaviours have been independently shown to be associated with adolescent and young adult smoking, drinking, and gambling. Nevertheless, the relation between early initiation into risky and addictive behaviours and CPs remains to be explored among pre-adolescents. Our study aims to examine the prospective relation between CPs in early primary school and pre-adolescent initiation into smoking, alcohol use, and gambling.

Method: Our study used data from participants in an ongoing prospective, longitudinal study at the Université de Sherbrooke to examine cigarette, alcohol, and gambling initiation among primary school-aged boys and girls with CPs. Children were recruited between the ages of 6 and 9 years from several low socioeconomic status public schools in diverse geographical regions of Quebec. Initiation into cigarettes, alcohol, and gambling was measured 1 year later.

Results: Children with CPs were found to be at greater risk for early initiation into smoking, alcohol, and gambling. These effects remained even once other known risk factors, such as poor parental supervision and child effortful control, were controlled for.

Conclusions: These results suggest that CPs present in early elementary school can predict early initiation in to potentially addictive behaviours among boys and girls. Implications for targeted preventive intervention are discussed.

Abrégé

Objectif : La participation des adolescents à des comportements à risques créant une dépendance, comme le tabagisme, l'utilisation de substances, et le jeu, a le potentiel de mener à une variété de problèmes sérieux. La présence de problèmes de conduite (PC) et un début précoce de comportements à risques créant une dépendance se sont indépendamment révélés être associés au tabagisme, à la consommation d'alcool et au jeu chez les adolescents et les jeunes adultes. Néanmoins, la relation entre le début précoce des comportements à risques créant une dépendance et les PC demeure à être explorée chez les pré-adolescents. Cette étude vise à examiner la relation prospective entre les PC au premier cycle du primaire et le début du tabagisme, de la consommation d'alcool et du jeu chez les pré-adolescents.

Méthode : La présente étude a utilisé les données des participants à une étude prospective, longitudinale en cours à l'Université de Sherbrooke pour examiner le début du tabagisme, de l'alcool et du jeu chez les garçons et filles d'âge scolaire au primaire ayant des PC. Les enfants ont été recrutés entre l'âge de 6 et 9 ans dans plusieurs écoles publiques de faible statut

¹ Université de Sherbrooke, Sherbrooke, Quebec

Corresponding Author:

Caroline E. Temcheff, PhD, Département de Psychoéducation, Université de Sherbrooke, 150, place Charles-LeMoyne, Suite 12830, Longueuil, QC J4K 0A8.
Email: caroline.temcheff@usherbrooke.ca

socio-économique (SSE), dans diverses régions géographiques du Québec. Le début du tabagisme, de l'alcool et du jeu a été mesuré un an plus tard.

Résultats : Les enfants ayant des PC étaient à risque accru d'un début précoce du tabagisme, de l'alcool et du jeu. Ces effets persistaient même après avoir contrôlé d'autres facteurs de risque comme une mauvaise supervision parentale et le contrôle volontaire de l'enfant.

Conclusions : Ces résultats suggèrent que les PC présents au premier cycle du primaire peuvent prédire un début précoce de comportements pouvant créer une dépendance chez les garçons et les filles. Les implications pour une intervention préventive ciblée sont discutées.

Keywords

longitudinal, conduct problems, gambling, substance abuse, early initiation

Adolescent participation in risky and addictive behaviours is a common phenomenon. Epidemiological research reveals that about 1 in 10 early adolescents and close to one-half of older adolescents report regular use of alcohol.¹ Regarding heavy episodic drinking, findings demonstrate that over one-quarter of students aged 15 to 17 years indicate having consumed 5 or more drinks in a row in the past month.² Data also suggest that about 3% to 11% report daily use of cigarettes.³ Further, prevalence research reveals approximately 9% to 19% of adolescents gamble for money on a frequent basis.⁴⁻⁶

While for many adolescents, participation in risk or addictive behaviours does not result in serious negative consequences, those who are initiated into these behaviours at young ages are at elevated risk for alcohol-related problems in adolescence,⁷⁻¹⁰ as well as nicotine dependence,¹¹ problematic alcohol consumption,¹² and at-risk gambling status¹³ in young adulthood. The fact that early onset of risk and addictive behaviours is related to various problematic outcomes draws attention to the importance of identifying the precursors to the acquisition and development of these behaviours among children and pre-adolescents. A better understanding of these precursors will ultimately assist in the development of preventive interventions aimed at problems before they occur.^{14,15}

Among the individual and familial risk factors identified in the literature as associated with youth addictive behaviours, a number are described as nonspecific¹⁴ given that they predict a wide range of youth maladaptive behaviours. Conduct problems (CPs) among children is widely recognized as an important nonspecific individual-level risk factor for addiction.¹⁶ Before examining predictors of early initiation into risky and addictive behaviours, we will explore the relation between CPs and addictive behaviours.

Childhood Conduct Problems and Risk or Addictive Behaviours

CPs refers to conduct disorder (CD) and oppositional defiant disorder (ODD), which are described in the Diagnostic and Statistical Manual of Mental Disorders (DSM) Fifth Edition,¹⁷ and are 2 mental health disorders that can co-occur.^{18,19} These mental disorders are manifested in behaviours that violate the rights of others, and (or) behaviours that bring a person into conflict

with societal norms. These disorders, as well as subclinical manifestations of these problems, will be referred to as CPs.

In a study designed to examine the relations between CPs, attention-deficit hyperactivity disorder (ADHD), and internalizing disorders at age 11 and substance use at age 14, King et al²⁰ observed that CPs were significantly associated with increased risk for regular use of alcohol and nicotine in early adolescence. King et al²⁰ found that CD and ODD increased the odds of regular use of nicotine and alcohol, while ADHD was not prospectively linked to regular use of alcohol. Another study²¹ revealed that elevations in the number of CP symptoms in early adolescence was prospectively associated with greater risk for membership in the early heavy drinking trajectory, which is characterized by an early onset and high frequency of binge drinking.

CPs have also been reported to be prospectively associated with the development of substance use disorders (SUDs) in young adulthood. A small number of longitudinal studies have found that CPs in adolescence significantly increase the risk for abuse of alcohol or dependence to tobacco in later adolescence and early adulthood.²²⁻²⁴ Conversely, attentional problems or ADHD are observed to be unrelated to increased risks of later alcohol abuse or nicotine dependence when CPs are controlled for.^{22,23}

A cross-sectional study²⁵ examining links between problem gambling and CD among adolescents and young adults reported a strong positive relation between the 2 problems. The relation was much stronger for those with an earlier age of onset of gambling problems (that is, age of onset of 14 years or younger) than those with a later age of onset (that is, age of onset of 15 years or older). Another investigation found that CD had one of the strongest associations with youth problem gambling, even after other concurrent problem behaviours (alcohol or marijuana problems or tobacco dependence) were taken into account.²⁶ However, no studies have examined prospective links between childhood CPs and pre-adolescent initiation into gambling.

Other Predictors of Pre-adolescent Initiation to Risk Taking and Addictive Behaviours

There has been little research attention devoted to the identification of childhood individual and familial risk factors

that contribute to pre-adolescent smoking, alcohol use, and gambling initiation. Within the extant literature, being male has been observed as a prospective predictor of smoking initiation during late childhood,^{27,28} as well as a prospective predictor of substance use initiation in general.²⁹ In addition, higher levels of impulsivity in kindergarten are shown to be predictive of self-reported gambling behaviour in late elementary school.^{30,31} Although not among pre-adolescents, there is also evidence for the developmental role of impulsivity as a risk factor for nicotine dependence,²² and gambling problems in adolescence.³²⁻³⁴ The related concept of low behavioural self-control has been identified to play a predictive role in cigarette smoking³⁵ and heavy drinking³⁶ in adolescence. Behavioural control is a component of the temperamental factor of effortful control, but effortful control also includes attentional control and the child's perceptual sensitivity.³⁷ Children with slower rates of increase of effortful control were more likely to use alcohol and other drugs in adolescence.³⁸

Data also reveal that familial characteristics play a significant role in the development of pre-adolescent risk and addictive behaviours. Parental use of tobacco is observed to be a robust antecedent risk factor for pre-adolescent smoking initiation.^{27,28,35,39} Likewise, parental alcohol use is shown to be a risk factor for pre-adolescent use of alcohol.^{10,40} Moreover, low levels of parental monitoring are reported to be a risk factor for general substance use initiation in pre-adolescence.⁴¹⁻⁴³

Our Study. Although early initiation and CPs have both been identified as independent risk factors for adolescent and young adult smoking, drinking, and gambling, the relation between these 2 risk variables remains unexplored. Few studies have investigated whether there are common or shared developmental antecedents for early initiation of gambling behaviour and other potentially addictive behaviours (smoking or alcohol consumption) among pre-adolescents. Our study aimed to evaluate the prospective association of childhood CPs and initiation of smoking, alcohol use, and gambling among pre-adolescent boys and girls, while controlling for the effects of other potential predictors.

Method

Participants

Participants were drawn from an ongoing longitudinal project from the Université de Sherbrooke in Quebec, on boys and girls referred to elementary school-based services for CPs. This project began in 2007-2008. Children in the study were French-speaking and were between the ages of 6.3 and 9.9 years at study inception ($N = 523$), with a mean age of 8.5 years. At time 2, children had a mean age of 9.5 years ($n = 501$, 4.2% attrition).

Participants were recruited from 8 public school boards in Quebec. The children with CPs were selected based on 2

criteria: obtain a score above the threshold of elevated risk on the DSM-Oriented Scales for CPs or for oppositional defiant problems (ODPs) of the Achenbach System of Empirically Based Assessment (ASEBA)⁴⁴ based on parent or teacher report (see the section on Predictor Variables), and were actively receiving specialized services for CPs in their school. All girls and an equal number of randomly selected boys who were less than 10 years of age and who met criteria were invited to participate. Children with an intellectual or physical disability or pervasive developmental disorder were excluded, as were children who were not living with at least one biological parent. The participation rate of families of children with CPs was 75.1%. Participants did not differ from nonparticipants in the proportion of girls, grade level, and the deprivation index of the school that the child attended. Children in the control group were selected from schools with a high deprivation index. The Ministry of Education of Quebec assigns schools a deprivation index that is established based on the family revenue, the education of the parents, and the employment status of the parents in the particular geographical region that the school serves.⁴⁵ The selection of the control group within schools with a high deprivation index was done to reduce school and residential neighborhood disparities with children in the CP group. Children in the control group had no CPs according to the teacher and parent-reported ASEBA, and did not receive any specialized services for CPs in their schools.

Our sample is comprised of participants who completed all questionnaires included in this study (at both times 1 and 2). In total, the sample included 176 boys with CPs, 116 girls with CPs, 98 matched control boys, and 100 matched control girls ($n = 490$).

Outcome Measures: Initiation into Smoking, Alcohol, and Gambling

A questionnaire from the National Longitudinal Survey of Children and Youth^{46,47} was used at time 2 to measure child smoking, alcohol, and gambling. Children were asked if they had ever tried: cigarettes (yes or no); drinking more than a few sips of alcohol (yes or no); and, gambling or betting on games of chance to gain money or other goods (yes or no). Overall, 35 children said that they had previously tried smoking, 131 had tried alcohol, and 53 had tried gambling at least once.

Predictor Variables

Conduct Problems. The presence of CPs was established using the DSM-Oriented Scales for CPs and ODPs of the parent and teacher report forms of ASEBA.⁴⁴ The DSM-Oriented Scales included items rated by a panel of experts as very consistent with symptoms of CD and ODD listed in the DSM-IV.⁴⁸ Items were scored on a 3-point scale ranging from 0 (not true) to 2 (very true or often true).

Family Poverty. Parents were asked to provide their family income. An income cutoff of Can\$25 000 was used to create this dichotomous variable.⁴⁹

Lack of Parental Supervision. A subscale of the Alabama Parenting Questionnaire⁵⁰ was used to measure parental monitoring. The scale consists of 10 items rated on a 5-point frequency scale ranging from 1 (never) to 5 (always).

Parental Antisocial Traits. A subscale of the Millon Clinical Multiaxial Inventory–III,⁵¹ constructed based on the DSM-IV⁴⁸ definition of antisocial personality disorder, is composed of 17 self-reported true or false items.

Parental Substance Abuse Problem. Parents were asked if either of the child's parents or parent's current romantic partners had a problem with alcohol consumption or with drug use since the birth of the child.

Effortful Control. Effortful control was measured using 4 subscales of the Children's Behavior Questionnaire–Short Form.⁵² The subscales, comprised of 26 items rated on 7-point scales ranging from 1 (extremely untrue) to 7 (extremely true), assessed attentional focusing, inhibitory control, low intensity pleasure, and perceptual sensitivity.

Procedure

Parents were presented with a full description of the study and signed a consent form, approved by the Institutional Ethics Board, for participation in the study and to obtain information about the child's behaviour from teachers. Child verbal assent was also obtained. Parents and children, accompanied by research assistants, completed questionnaires separately in their home. The teacher report was obtained by telephone. Parents, children, and teachers received financial compensation for participating. Interviews were performed by graduate-level research assistants, all of whom underwent a formal 3-day training session.

Statistical Analysis

To address the research goal of our study, logistic regression was selected as the data analytic method given that the outcome variable was dichotomous (previous initiation into smoking, alcohol use, or gambling, compared with no previous initiation) and that the distributions of the independent and control variables (CPs, poverty, and parental antisocial behaviour) were not likely to satisfy the assumptions of normality.⁵³ The logistic regression was undertaken to evaluate the contribution of each predictor while controlling for the effects of the other predictor variables.⁵³ Once all main effects were examined, further regression analyses were carried out examining interactions between CPs and each of the predictors. All were found nonsignificant in the prediction of any of the outcomes.

Results

Correlations between predictor variables and outcomes are shown in Table 1.

Predicting Early Initiation into Cigarette Use

A logistic regression analysis was performed to determine whether the presence of CPs in childhood predicted early initiation into cigarette use. A test of the model against the intercept-only model was statistically significant ($\chi^2 = 33.50$, $df = 8$, $P < 0.001$). Overall, the regression equation accounted for 16% of the variance in cigarette initiation among children in our sample.

The contribution of each of the individual predictors on cigarette initiation, while controlling for the other variables in the model, is summarized in Table 2. The presence of a child with CPs substantially increased the likelihood of initiation into cigarettes ($\beta = 2.13$, $P < 0.01$). Children with CPs were found to be over 8 times more likely to have tried cigarettes by about age 9 than their counterparts without CPs. This effect was present after the effects of sex, age, poverty, lack of parental supervision, parental antisocial behaviour and substance use problems, and child effortful control were accounted for. None of the control variables were significant predictors of cigarette initiation once the effects of CPs were accounted for.

Predicting Early Initiation into Alcohol Use

A logistic regression analysis was performed to determine whether the presence of CPs in childhood predicted early initiation into alcohol consumption. A test of the model against the intercept-only model was statistically significant ($\chi^2 = 19.81$, $df = 8$, $P < 0.01$). Overall, the model accounted for 6% of the variance in alcohol initiation among children in our sample.

The contribution of each of the individual predictors on alcohol initiation, while controlling for the other variables in the model, is summarized in Table 3. The presence of a CP was found to be a significant predictor of early initiation into alcohol consumption ($\beta = 0.53$, $P < 0.05$), with children with CPs being 1.7 times more likely to have been initiated into alcohol use than their peers without CPs.

The presence of substance abuse problems in at least one parent was found to be a significant negative predictor of alcohol initiation for children ($\beta = -0.64$, $P < 0.05$), suggesting that those with at least one parent with a substance abuse problem were less likely to have tried alcohol than those without parents with substance abuse problems.

Predicting Early Gambling Involvement

A logistic regression analysis was performed to determine whether the presence of CPs in childhood predicted early initiation into gambling. A test of the model against the intercept-only model was statistically significant ($\chi^2 =$

Table 1. Correlations between predictor variables and outcome variables.

	1. Cigarette initiation	2. Alcohol initiation	3. Gambling initiation	4. Conduct problems	5. Sex	6. Age	7. Poverty	8. Lack of parental supervision	9. Parental antisocial behaviour	10. Parental substance use problem	11. Child effortful control
1.	—	0.21 ^a	0.06	0.19 ^a	-0.01	0.09 ^b	0.05	0.13 ^a	0.12 ^a	0.15 ^a	-0.13 ^a
2.		—	0.13 ^a	0.11 ^b	-0.04	0.10 ^b	-0.05	0.04	0.06	-0.07	-0.08
3.			—	0.15 ^a	-0.11 ^b	-0.01	0.06	0.06	0.08	0.00	-0.12 ^a
4.				—	-0.11 ^b	0.02	0.15 ^a	0.27 ^a	0.31 ^a	0.26 ^a	-0.55 ^a
5.					—	-0.01	0.06	0.06	0.05	0.02	0.21 ^a
6.						—	-0.03	0.08	0.01	0.03	-0.09 ^b
7.							—	0.10 ^b	0.26 ^a	0.21 ^a	-0.14 ^a
8.								—	0.27 ^a	0.19 ^a	-0.24 ^a
9.									—	0.31 ^a	-0.26 ^a
10.										—	-0.21 ^a
11.											—

^a $p < 0.01$; ^b $p < 0.05$.

Table 2. Logistic regression analysis predicting cigarette initiation (*n* = 490).

Predictor	β	SE	<i>P</i>	OR	95% CI
Sex	-0.01	0.39	0.99	0.99	0.47 to 2.12
Age	0.42 ^a	0.21	0.05	1.52	1.00 to 2.31
Poverty	0.01	0.42	0.99	1.00	0.44 to 2.30
Lack of parental supervision	0.59	0.57	0.30	1.8	0.59 to 5.55
Parental antisocial behaviour	0.04	0.08	0.60	1.04	0.90 to 1.21
Child effortful control	-0.09	0.30	0.77	0.92	0.51 to 1.65
Parental substance use problem	0.64	0.39	0.10	1.89	0.88 to 4.07
Child conduct problems	2.13^b	0.78	0.01^c	8.42	1.81 to 39.09

Nagelkerke R^2 = 0.16.^a P < 0.10; ^b P < 0.01; ^c P = 0.007. β = parameter estimate; OR = odds ratio; SE = standard error.**Table 3.** Logistic regression analysis predicting alcohol initiation (*n* = 490).

Predictor	β	SE	<i>P</i>	OR	95% CI
Sex	-0.07	0.22	0.75	0.93	0.61 to 1.43
Age	0.22 ^a	0.11	0.05	1.25	0.99 to 1.56
Poverty	-0.40	0.28	0.15	0.67	0.39 to 1.16
Lack of parental supervision	0.06	0.41	0.89	1.06	0.47 to 2.37
Parental antisocial behaviour	0.08	0.05	0.11	1.09	0.98 to 1.20
Child effortful control	-0.08	0.18	0.67	0.93	0.66 to 1.31
Parental substance use problem	-0.64 ^b	0.28	0.02	0.53	0.30 to 0.91
Child conduct problems	0.53^b	0.27	0.05^c	1.70	1.01 to 2.86

Nagelkerke R^2 = 0.06.^a P < 0.10; ^b P < 0.05; ^c P = 0.048. β = parameter estimate; OR = odds ratio; SE = standard error.

21.12, *df* = 8, P < 0.01). Overall, the model accounted for 9% of the variance in gambling initiation among children in our sample.

The contribution of each of the individual predictors on gambling initiation, while controlling for the other variables in the model, is summarized in Table 4. Consistent with other literature, sex emerged as a statistically significant predictor, with the odds of a girl having been initiated into gambling being about one-half that of boys (β = -0.74, P < 0.05). In addition, having CPs was directly predictive of gambling initiation among our child participants, even when the effects of sex, poverty, poor parental supervision, parental antisocial, and substance use problems were accounted for (β = 1.00, P < 0.05). None of the control variables were statistically significant predictors of gambling initiation once the effects of sex and presence of CPs were accounted for.

Table 4. Logistic regression analysis predicting gambling initiation (*n* = 490).

Predictor	β	SE	<i>P</i>	OR	95% CI
Sex	-0.74 ^a	0.34	0.03	0.48	0.25 to 0.92
Age	-0.05	0.16	0.73	0.95	0.70 to 1.29
Poverty	0.34	0.35	0.33	1.41	0.70 to 2.82
Lack of parental supervision	0.37	0.54	0.49	1.45	0.50 to 4.17
Parental antisocial behaviour	0.06	0.07	0.43	1.06	0.92 to 1.21
Child effortful control	-0.07	0.25	0.78	0.93	0.58 to 1.51
Parental substance use problem	-0.49	0.38	0.20	0.62	0.29 to 1.29
Child conduct problems	1.00^a	0.42	0.02	2.71	1.19 to 6.18

Nagelkerke R^2 = 0.09.^a P < 0.05. β = parameter estimate; OR = odds ratio; SE = standard error.

Discussion

These results suggest that even at about 9.5 years of age, children with CPs were over 1.5 times more likely to have been initiated into alcohol use, over twice as likely to have been initiated into gambling, and over 8 times more likely to have been initiated into cigarette use than their peers without CPs. Further, these effects were present even after controlling for sex and age, as well as other variables, such as parental monitoring and child effortful control.

Our study found a lack of predictive significance of child effortful control on early initiation into cigarettes, alcohol, and gambling. Although childhood impulsivity and low behavioural control have been shown to be a risk factor for early initiation into addictive behaviours,³⁰ no study has yet to examine the effects of childhood effortful control and CPs simultaneously in the prediction of initiation into cigarettes, alcohol, and gambling in middle elementary school. We believe that childhood effortful control is a more complete measure of a child's self-regulation, as it includes not only behavioural control, but also attentional control.

It appears from our findings that, at least among the young disadvantaged sample used in our study, having CPs is a more robust predictor of very early initiation to cigarettes, alcohol, and gambling, than is child effortful control. Although the reason for the predictive ability of childhood CPs over and above effortful control cannot be ascertained given the current methodology, these results are in line with previous research^{20,22,23} suggesting that CPs might be more strongly associated with initiation into risky and addictive behaviours, than ADHD. One possible reason could be that with respect to very early initiation into substances and gambling, early experimentation has more to do with social milieus and rule breaking, than with a lack of effortful control.

Research^{38,54-59} has previously demonstrated the heightened risk for alcohol abuse and alcohol-related problems

among adolescent children of alcoholics. However, the results of our study suggest a negative relation between parental substance abuse problems and child early initiation into alcohol. Specifically, children with biological parents who suffer from substance abuse problems were almost one-half as likely to have been initiated into alcohol as their peers whose parents did not have substance use problems. While this finding may initially seem inconsistent with the literature, most previous research has focused on alcohol consumption and alcohol problems among adolescents, and not early initiation among elementary school-aged children. It is important to note that the variable was also recoded and tested to reflect only those children who currently lived with the parent with the substance abuse problem. Regardless of whether the child lived with the parent with the substance abuse problem or not, the negative relation between parental SUD and child early initiation remained the same. This suggests that the negative relation is not due to the absence of the parent with the substance abuse problem, leaving the child with the other, unaffected parent.

There are several possible explanations to this finding. First, it is possible that young children who have alcoholic parents might have more experience with the negative aspects of alcohol misuse and therefore may be less interested in trying it for themselves in these early years. Second, when one biological parent suffers from a SUD, the other parent might be more vigilant about not allowing children access to alcohol at early ages. Whatever the reason for this negative relation, it is evident from past studies that, somewhere between childhood and early-to mid-adolescence, parental substance problems become a significant and positive predictor of adolescent substance misuse and problems. The timing and mechanisms of the change in the direction of the relation between parental substance abuse problems and child or adolescent substance use or misuse are unclear; however, continued follow-up of this cohort of young children with CPs may help shed some light on the associated processes in coming years.

Strengths and Limitations

Longitudinal designs employ a prospective approach instead of relying on retrospective data, which is subject to recall bias. Causal inferences can be drawn more credibly in prospective longitudinal studies because of the establishment of time sequencing in the prediction of outcomes. However, in this case as in most cases, conclusions remain speculative as a function of the methodology, which is correlational. Replication of results across studies, using different samples is a necessary step in confirming the generality of longitudinal results.⁶⁰

Another strength of our design lies in the very young age of respondents. The goal of our study was to investigate very early initiation into potentially addictive behaviours, such as

cigarette and alcohol use and gambling among children. However, given the young age at which the outcome data was collected, there was only about 12 months of time elapsed between the collection of the predictors and the outcomes. Replication of results in future years with these children will help clarify the picture in terms of the development of addictive behaviours.

While the effect size of the presence of CPs on cigarette initiation was substantial, the effect sizes in the prediction of alcohol and gambling initiation were modest. Nevertheless, considering the very young age of participants at the time of outcome data collection, the results offer new information as to the early predictors of substance use and gambling initiation, as well as suggesting possible avenues for preventative intervention in childhood.

Conclusions

Knowing how to identify and help children in elementary school who might be at high risk for early initiation into risky and addictive behaviours might help mitigate a host of negative long-term consequences. The fact that some of the children with CPs are at an increased risk for becoming initiated into cigarettes and gambling at very early ages, even after controlling for sex, poor parental supervision, and other risk factors, has significant implications for prevention. These results suggest that the elementary school years may be an appropriate time to engage in prevention programs for substance use and gambling among children with CPs. Further, given that these children are at heightened risk for early initiation into gambling also suggests that prevention of gambling should be considered alongside prevention of other risky and addictive behaviours when designing comprehensive prevention programs for children with CPs.

Acknowledgements

The authors thank France Girard, Annie Lemieux, and the entire Université de Sherbrooke Longitudinal Project team for their assistance in data collection and analysis. Finally, we are most indebted to the participants of the study.

Declaration of Conflicting Interests

The author(s) declare no potential conflicts of interest regarding the research, authorship, and (or) publication of this article.

Funding

The author(s) declared receipt of the following financial support for the research, authorship, and/or publication of this article: This research was partially supported by grants from the Canadian Institutes for Health Research (funding reference number 82694), the Social Sciences and Humanities Research Council (grant 37890), and the Fonds québécois de la recherche sur la société et la culture (funding reference number 2013-NP-165564).

References

- Swendsen J, Burstein M, Case B, et al. Use and abuse of alcohol and illicit drugs in US adolescents: results of the National Comorbidity Survey-Adolescent Supplement. *Arch Gen Psychiatry*. 2012;69(4):390-398.
- Miller JW, Naimi TS, Brewer RD, et al. Binge drinking and associated health risk behaviors among high school students. *Pediatrics*. 2007;119(1):76-85.
- Johnston LD, O'Malley PM, Bachman JG, et al. Monitoring the Future national survey results on drug use, 1975-2010: volume I, secondary school students. Ann Arbor (MI): Institute for Social Research, University of Michigan; 2011.
- Volberg RA, Gupta R, Griffiths MD, et al. An international perspective on youth gambling prevalence studies. *Int J Adolesc Med Health*. 2010;22(1):3-38.
- Welte JW, Barnes GM, Tidwell MC, et al. The prevalence of problem gambling among US adolescents and young adults: results from a national survey. *J Gambl Stud*. 2008;24(2):119-133.
- Welte JW, Barnes GM, Tidwell MC, et al. Gambling and problem gambling across the lifespan. *J Gambl Stud*. 2011;27(1):89-99.
- Ellickson PI, Tucker JS, Klein DJ, et al. Prospective risk factors for alcohol misuse in late adolescence. *J Stud Alcohol*. 2001;62(6):773-782.
- Fergusson DM, Lynskey MT, Horwood LJ. Childhood exposure to alcohol and adolescent drinking patterns. *Addiction*. 1994;89(8):1007-1016.
- Gruber E, DiClemente RJ, Anderson MM, et al. Early drinking onset and its association with alcohol use and problem behavior in late adolescence. *Prev Med*. 1996;25(3):293-300.
- Hawkins JD, Graham JW, Maguin E, et al. Exploring the effects of age of alcohol use initiation and psychosocial risk factors on subsequent alcohol misuse. *J Stud Alcohol*. 1997;58(3):280-290.
- Behrendt S, Wittchen HU, Hofler M, et al. Transitions from first substance use to substance use disorders in adolescence: is early onset associated with a rapid escalation? *Drug Alcohol Depend*. 2009;99(1-3):68-78.
- Buchmann AF, Schmid B, Blomeyer D, et al. Impact of age at first drink on vulnerability to alcohol-related problems: testing the marker hypothesis in a prospective study of young adults. *J Psychiatr Res*. 2009;43(15):1205-1212.
- Winters KC, Stinchfield RD, Botzet A, et al. A prospective study of youth gambling behaviors. *Psychol Addict Behav*. 2002;16(1):3-9.
- Zucker RA, Donovan JE, Masten AS, et al. Early developmental processes and the continuity of risk for underage drinking and problem drinking. *Pediatrics*. 2008;121(Suppl 4): S252-S272.
- Shead NW, Derevensky JL, Gupta R. Risk and protective factors associated with youth problem gambling. *Int J Adolesc Med Health*. 2010;22(1):39-58.
- Tarter RE, Kirisci L, Habeych M, et al. Neurobehavior disinhibition in childhood predisposes boys to substance use disorder by young adulthood: direct and mediated etiologic pathways. *Drug Alcohol Depend*. 2004;73(2):121-132.
- American Psychiatric Association (APA). Diagnostic and statistical manual of mental disorders. 5th ed. Washington (DC): APA; 2013.
- Maughan B, Rowe R, Messer J, et al. Conduct disorder and oppositional defiant disorder in a national sample: developmental epidemiology. *J Child Psychol Psychiatry*. 2004;45:609-621.
- Costello E, Mustillo S, Erkanli A, et al. Prevalence and development of psychiatric disorders in childhood and adolescence. *Arch Gen Psychiatry*. 2003;60:837-844.
- King SM, Iacono WG, McGue M. Childhood externalizing and internalizing psychopathology in the prediction of early substance use. *Addiction*. 2004;99(12):1548-1559.
- Chassin L, Pitts SC, Prost J. Binge drinking trajectories from adolescence to emerging adulthood in a high-risk sample: predictors and substance abuse outcomes. *J Consult Clin Psychol*. 2002;70(1):67-78.
- Elkins IJ, McGue M, Iacono WG. Prospective effects of attention-deficit/hyperactivity disorder, conduct disorder, and sex on adolescent substance use and abuse. *Arch Gen Psychiatry*. 2007;64(10):1145-1152.
- Fergusson DM, Horwood LJ, Ridder EM. Conduct and attentional problems in childhood and adolescence and later substance use, abuse and dependence: results from a 25-year longitudinal study. *Drug Alcohol Depend*. 2007;88(Suppl 1):S14-S26.
- Sung M, Erkanli A, Angold A, et al. Effects of age at first substance use and psychiatric comorbidity on the development of substance use disorders. *Drug Alcohol Depend*. 2004;75(3):287-299.
- Welte JW, Barnes GM, Tidwell MCO, et al. Association between problem gambling and conduct disorder in a national survey of adolescents and young adults in the United States. *J Adolesc Health*. 2009;45(4):49-61.
- Barnes GM, Welte JW, Hoffman JH, et al. The co-occurrence of gambling with substance use and conduct disorder among youth in the United States. *Am J Addiction*. 2011;20(2):166-173.
- Milton B, Cook PA, Dugdill L, et al. Why do primary school children smoke? A longitudinal analysis of predictors of smoking uptake during pre-adolescence. *Public Health*. 2004;118(4):247-255.
- O'Loughlin J, Paradis G, Renaud L, et al. One-year predictors of smoking initiation and of continued smoking among elementary schoolchildren in multiethnic, low-income, inner-city neighbourhoods. *Tob Control*. 1998;7(3):268-275.
- Kaplow JB, Curran PJ, Dodge KA, et al. Child, parent and peer predictors of early-onset substance use: a multisite longitudinal study. *J Abnorm Child Psychol*. 2002;30(3):199-216.
- Pagani LS, Derevensky JL, Japel C. Predicting gambling behavior in sixth grade from kindergarten impulsivity. *Arch Pediatr Adol Med*. 2009;163(3):238-243.
- Vitaro F, Wanner B. Predicting early gambling in children. *Psychol Addict Behav*. 2011;25(1):118-126.
- Dussault F, Brendgen M, Vitaro F, et al. Longitudinal links between impulsivity, gambling problems and depressive

- symptoms: a transactional model from adolescence to early adulthood. *J Child Psychol Psychiatry*. 2011;52(2):130-138.
33. Vitaro F, Arseneault L, Tremblay RE. Impulsivity predicts problem gambling in low SES adolescent males. *Addiction*. 1999;94(4):565-575.
 34. Vitaro F, Wanner B, Ladouceur R, et al. Trajectories of gambling during adolescence. *J Gambl Stud*. 2004;20(1):47-69.
 35. Jackson C, Henriksen L, Dickinson D, et al. A longitudinal study predicting patterns of cigarette smoking in late childhood. *Health Educ Behav*. 1998;25(4):436-447.
 36. Griffin KW, Botvin GJ, Epstein JA, et al. Psychosocial and behavioral factors in early adolescence as predictors of heavy drinking among high school seniors. *J Stud Alcohol*. 2000; 61(4):603-606.
 37. Rothbart MK. *Becoming who we are: temperament and personality in development*. New York (NY): Guilford Press; 2011. p 324.
 38. Wong MM, Nigg JT, Zucker RA, et al. Behavioral control and resiliency in the onset of alcohol and illicit drug use: a prospective study from preschool to adolescence. *Child Dev*. 2006; 77(4):1016-1033.
 39. Fleming CB, Kim H, Harachi TW, et al. Family processes for children in elementary school as predictors of smoking initiation. *J Adolescent Health*. 2002;30(3):184-189.
 40. Hung CC, Yen LL, Wu WC. Association of parents' alcohol use and family interaction with the initiation of alcohol use by sixth graders: a preliminary study in Taiwan. *BMC Public Health*. 2009;9:172. doi: 10.1186/1471-2458-9-172.
 41. Chilcoat HD, Anthony JC. Impact of parent monitoring on initiation of drug use through late childhood. *J Am Acad Child Adolesc Psychiatry*. 1996;35(1):91-100.
 42. Chilcoat HD, Breslau N. Pathways from ADHD to early drug use. *J Am Acad Child Adolesc Psychiatry*. 1999;38(11):1347-1354.
 43. Chilcoat HD, Dishion TJ, Anthony JC. Parent monitoring and the incidence of drug sampling in urban elementary school children. *Am J Epidemiol*. 1995;141(1):25-31.
 44. Achenbach TM, Rescorla LA. *Manual for the ASEBA school-age forms & profiles*. Burlington (VT): University of Vermont, Research Center for Children, Youth, and Families; 2001. p 238.
 45. Gouvernement du Québec. *Atlas de la défavorisation*. Québec (QC): Gouvernement du Québec; 2012.
 46. Statistics Canada, Human Resources Development Canada. *Enquête longitudinale nationale sur les enfants et les jeunes: matériel d'enquête pour la collecte des données de 1996-1997, cycle 2*. Catalogue no: 89FOO77XIF. Ottawa (ON): Government of Canada; 1999.
 47. Statistics Canada, Human Resources Development Canada. *National Longitudinal Survey of Children and Youth: survey instruments for 1996-1997 data collection, cycle 2*. Catalogue no: 89FOO77XPE. Ottawa (ON): Government of Canada; 1999.
 48. American Psychiatric Association (APA). *Diagnostic and statistical manual of mental disorders*. 4th ed. Washington (DC): APA; 1994. p 886.
 49. Institut de la statistique du Québec. *Seuils du faible revenu, Mesure du Faible Revenu (MFR)-seuils, avant impôt, selon la taille du ménage, Québec, 1996-2011* [Internet]. Québec (QC): Gouvernement du Québec; 2014 [cited 2014 Sep 22]. Available from: http://www.stat.gouv.qc.ca/statistiques/conditions-vie-societe/revenu/faible-revenu/seuils_mfr_qcavi.htm. French.
 50. Shelton KK, Frick PJ, Wootton J. Assessment of parenting practices in families of elementary school-age children. *J Clin Child Psychol*. 1996;25(3):317-329.
 51. Millon T, Davis RD, Millon C. *Manual for the Millon Clinical Multiaxial Inventory—III*. 2nd ed. Minneapolis (MN): National Computer Systems; 1997.
 52. Putnam SP, Rothbart MK. Development of short and very short forms of the Children's Behavior Questionnaire. *J Pers Assess*. 2006;87(1):102-112.
 53. Tabachnick BG, Fidell LS. *Using multivariate statistics*. 6th ed. Boston (MA): Allyn and Bacon; 2001. p 1024.
 54. Ary DV, Tildesley E, Hops H, et al. The influence of parent, sibling, and peer modeling and attitudes on adolescent use of alcohol. *Subst Use Misuse*. 1993;28(9):853-880.
 55. Engels RCME, Knibbe RA, De Vries H, et al. Influences of parental and best friends' smoking and drinking on adolescent use: a longitudinal study. *J Appl Soc Psychol*. 1999;29(2): 337-361.
 56. Getz JG, Bray JH. Predicting heavy alcohol use among adolescents. *Am J Orthopsychiatry*. 2005;75(1):102-116.
 57. Latendresse SJ, Rose RJ, Viken RJ, et al. Parenting mechanisms in links between parents' and adolescents' alcohol use behaviors. *Alcohol Clin Exp Res*. 2008;32(2): 322-330.
 58. Seljamo S, Aromaa M, Koivusilta L, et al. Alcohol use in families: a 15-year prospective follow-up study. *Addiction*. 2006; 101(7):984-992.
 59. van der Zwaluw CS, Scholte RHJ, Vermulst AA, et al. Parental problem drinking, parenting, and adolescent alcohol use. *J Behav Med*. 2008;31(3):189-200.
 60. Serbin L, Karp J. Intergenerational studies of parenting and the transfer of risk from parent to child. *Curr Dir Psychol Sci*. 2003;12(4):138-142.